

# Long cycles in critical graphs

Noga Alon \*

Michael Krivelevich †

Paul Seymour ‡

## Abstract

It is shown that any  $k$ -critical graph with  $n$  vertices contains a cycle of length at least  $2\sqrt{\log(n-1)/\log(k-2)}$ , improving a previous estimate of Kelly and Kelly obtained in 1954.

## 1 Introduction

A graph is  $k$ -critical if its chromatic number is  $k$  but the chromatic number of any proper subgraph of it is at most  $k-1$ . For a graph  $G$ , let  $L(G)$  denote the maximum length of a cycle in  $G$ , and define  $L_k(n) = \min L(G)$  where the minimum is taken over all  $k$ -critical graphs  $G$  with at least  $n$  vertices. Answering a problem of Dirac, Kelly and Kelly [3] proved that for every fixed  $k > 2$  the function  $L_k(n)$  tends to infinity as  $n$  tends to infinity. They also showed that  $L_4(n) \leq O(\log^2 n)$ , and after several improvements by Dirac and Read, Gallai [2] proved that for every fixed  $k \geq 4$  there are infinitely many values of  $n$  for which

$$L_k(n) \leq \frac{2(k-1)}{\log(k-2)} \log n.$$

This is the best known upper bound for  $L_k(n)$ . The best known lower bound, proved in [3], is that for every fixed  $k \geq 4$  there is some  $n_0(k)$  such that for all  $n > n_0(k)$

$$L_k(n) \geq \left( \frac{(2 + o(1)) \log \log n}{\log \log \log n} \right)^{1/2}, \quad (1)$$

where the  $o(1)$  term tends to 0 as  $n$  tends to infinity.

Note that the gap between the upper and lower bounds given above is exponential for fixed  $k$ , and the problem of determining the asymptotic behaviour of  $L_k(n)$  more accurately is still open; see also [1], Problem 5.11 for some additional relevant references.

---

\*School of Mathematics, Institute for Advanced Study, Princeton, NJ 08540, and Raymond and Beverly Sackler Faculty of Exact Sciences, Tel Aviv University, Tel Aviv, Israel. Email: noga@math.tau.ac.il. Research supported in part by a Sloan Foundation grant 96-6-2 and by a State of New Jersey grant.

†DIMACS Center, Rutgers University, Piscataway, NJ 08854, USA. Email: mkrivele@dimacs.rutgers.edu. Research supported by a DIMACS Postdoctoral Fellowship.

‡Department of Mathematics, Princeton University, Princeton, NJ 08544. Email: pds@math.princeton.edu. Research supported in part by ONR grant N00014-97-1-0512.

In the present note we improve the lower bound given in (1) and show that in fact  $L_k(n) \geq \Omega(\sqrt{\log n / \log(k-1)})$  for every  $n$  and  $k \geq 4$ . (Note that trivially  $L_3(n) = n$ .) The precise result we prove is the following.

**Theorem 1** *Let  $G$  be a  $k$ -critical graph on  $n$  vertices, and let  $t$  denote the length of the longest path in it. Then*

$$n \leq 1 + \sum_{j=0}^{t-1} s(j, k) \quad (2)$$

where

$$s(j, k) = j! \text{ for } j \leq k-3 \text{ and } s(j, k) = (k-2)!(k-2)^{j-k+2} \text{ for } j \geq k-2. \quad (3)$$

Therefore, any  $k$ -critical graph on  $n$  vertices contains a path of length at least  $\log(n-1)/\log(k-2)$  and a cycle of length at least  $2\sqrt{\log(n-1)/\log(k-2)}$ .

We note that the construction of Gallai shows that there are infinitely many values of  $n$  for which there is a  $k$ -critical graph on  $n$  vertices with no path of length greater than  $\frac{2(k-1)}{\log(k-2)} \log n$ , showing that the statement of the above theorem for paths is nearly tight for fixed  $k$ .

## 2 The Proof

Suppose  $k \geq 4$ , and let  $G = (V, E)$  be a  $k$ -critical graph on  $n$  vertices. Fix  $v_0 \in V$ , and let  $T$  be a depth first search (= DFS) spanning tree of  $G$  rooted at  $v_0$ . Denote the *depth* of  $T$ , (that is, the maximum length of a path from  $v_0$  to a leaf) by  $r$ , and recall that all non-tree edges of  $G$  are backward edges, that is, they connect a vertex of  $T$  with some ancestor of it in the tree. Call an edge  $uv$  of  $T$ , where  $u$  is the parent of  $v$ , an edge of *type*  $j$ , if the unique path in  $T$  from  $v_0$  to  $u$  has length  $j$ . Note that the type of each edge is an integer between 0 and  $r-1$ .

**Claim:** The number of edges of type  $j$  in  $T$  is at most  $s(j, k)$ , where  $s(j, k)$  is given in (3).

**Proof:** Assign to each edge  $e = uv$  of type  $j$  in  $T$ , where  $u$  is the parent of  $v$ , a word  $S_e$  of length  $j+1$  over the alphabet  $K = \{0, 1, 2, \dots, k-2\}$  as follows. Let  $v_0, v_1, \dots, v_j = u$  be the unique path in  $T$  from the root  $v_0$  to  $u$ . Let  $F_e$  be a proper coloring of  $G - e$  by the  $k-1$  colors in  $K$  such that  $F_e(v_i) \leq i$  for all  $i \leq k-2$ . Then  $S_e = (F_e(v_0), F_e(v_1), \dots, F_e(v_j))$ . The crucial observation is the fact that if  $e$  and  $e'$  are distinct tree edges of type  $j$ , then  $S_e \neq S_{e'}$ . Indeed, let  $e = uv$  be as above and suppose  $e' = u'v'$  is another edge of type  $j$ , where  $u'$  is the parent of  $v'$ . Let  $w$  be the lowest common ancestor of  $u$  and  $u'$  (which may be  $u$  itself, if  $u = u'$ ), and suppose  $S_e = S_{e'}$ . Then the two colorings  $F_e$  and  $F_{e'}$  coincide on the tree path from  $v_0$  to  $w$ . Let  $y$  be the vertex following  $w$  on the tree-path from  $v_0$  to  $v$  and let  $T_y$  be the subtree of  $T$  rooted at  $y$ . Define a coloring  $H$  of  $G$  as follows; for each vertex  $z$  of  $G$ ,  $H(z) = F_e(z)$  if  $z \notin T_y$ , and  $H(z) = F_{e'}(z)$  if  $z \in T_y$ . It is easy to check that since the only edges of  $G$  connecting  $T_y$  with the rest of the graph lead from  $T_y$  to the path from  $v_0$  to  $w$ , the coloring  $H$  is a proper coloring of  $G$  with  $k-1$  colors. This contradicts the

assumption that the chromatic number of  $G$  is  $k$ , and hence proves the required fact. Since every word  $S_e$  corresponds to a proper coloring of a path of length  $j+1$  in which the color of vertex number  $i$  is at most  $i$  (for all  $0 \leq i \leq j$ ), the number of possible distinct words is at most  $j!$  for  $j \leq k-3$ , and at most  $(k-2)!(k-2)^{j-k+2}$  if  $j \geq k-2$ . This completes the proof of the Claim.

By the above claim, the total number,  $n-1$ , of edges of  $T$  satisfies  $n-1 \leq \sum_{j=0}^{r-1} s(j, k)$ . Since  $r$  is the depth of the tree,  $G$  contains a path of length  $r$ , showing that  $t \geq r$  and hence implying (2). As  $k \geq 4$ , the right-hand-side of (2) is easily checked to be at most  $1 + (k-2)^{t-1}$ , implying that the maximum length of a path in  $G$  is at least  $\log(n-1)/\log(k-2)$ . Since  $G$  is 2-connected, it follows, by a theorem of Dirac (cf., e.g., [4]), that it contains a cycle of length at least  $2\sqrt{t}$ , completing the proof.  $\square$

**Remark 1.** It is easy to check that the above theorem implies that if  $k \geq 4$  then any  $k$ -critical graph  $G$  on  $n$  vertices contains an odd cycle of length at least  $\sqrt{\log(n-1)/\log(k-2)}$ . Indeed, let  $C$  be a longest cycle in  $G$ . If it is odd, the desired result follows, by Theorem 1. Otherwise, let  $A$  be an odd cycle in  $G$ . If  $A$  and  $C$  are vertex disjoint, there are, by the 2-connectivity of  $G$ , two internally disjoint paths from  $A$  to  $C$  providing an odd cycle containing at least half of  $C$ . A similar argument gives the same conclusion if  $A$  and  $C$  share only one common vertex. If they have more common vertices, split the edges of  $A$  not in  $C$  into paths that intersect  $C$  only in their ends. Then, there is such a path whose union with  $C$  is not 2-colorable (as otherwise the union of  $A$  and  $C$  would have been 2-colorable). Thus, in this case too we obtain an odd cycle containing at least half of  $C$ , and the required result follows from Theorem 1. Note that this shows that any large  $k$ -critical graph contains a large 3-critical subgraph. The problem of deciding if every large  $k$ -critical graph contains a large  $s$  critical graph for other values of  $k > s \geq 3$ , which is mentioned in [1], Problem 5.6, remains open.

**Remark 2.** A very simple proof of the fact that any 2-connected graph  $G$  containing a path  $P$  of length at least  $2s^2$  contains a cycle of length at least  $2s$  is as follows. If the distance in  $G$  between the two ends  $x$  and  $y$  of the path is at least  $s$ , then the union of two internally disjoint paths between  $x$  and  $y$  forms a cycle of length at least  $2s$ . Otherwise, consider a shortest path between  $x$  and  $y$ , and list its intersection points with the path  $P$ . Then the distance along  $P$  between some two such consecutive intersection points must be at least  $2s^2/s = 2s$ , providing, again, the required cycle. Although the proof in [4] gives a slightly better constant, the above argument is much simpler.

## References

- [1] T. Jensen and B. Toft, **Graph Coloring Problems**, Wiley, New York, 1995.
- [2] T. Gallai, Kritische Graphen I, Publ. Math. Inst. Hungar. Acad. Sci. 8 (1963), 165-192.

- [3] J. B. Kelly and L. M. Kelly, Paths and circuits in critical graphs, *Amer. J. Math.* **76** (1954), 786-792.
- [4] L. Lovász, **Combinatorial Problems and Exercises**, North Holland, Amsterdam, 1979, Problem 10.29.